Mantoux test koże za tuberkulozu

Developed by the
Minnesota Department of Health,
Tuberculosis Prevention and Control Program
August 2002 phone: (612) 676-5414

This is one in a series of four patient education pamphlets that addresses TB to be used in various clinical situations.

Mantoux Skin Test for Tuberculosis, Bosnian/Croatian/Serbian

Tuberkuloza i tuberkulozna infekcija

TB se mo`e izlije~iti uzimanjem lijekova

Tuberkuloza je ozbiljna bolest koju uzrokuju bakcili koji se razmno`avaju u tijelu i o{te}uju tkivo. Naj~e{}e utje~e na plu}a. TB se prenosi sa jedne osobe na drugu osobu zrakom. TB se mo`e izlije~iti lijekovima.

Znakovi i simptomi **TB** obuhva}aju:

- jaki ka{alj
- iska{ljavanje ispljuvka
- iska{ljavanje krvi
- bol u grudima kad ka{ljete
- gubitak te`ine
- povi{ena temperatura
- poja~ano znojenje tokom no}i
- bol zara`enog dijela tijela

Kad osoba koja je zara`ena sa TB na plu}ima ka{lje, ki{e ili vi~e TB bakcili se mogu rasprskati u zrak. Ostale osobe mogu udisati TB bakcile u plu}a. To se zove TB infekcija.

Osobe sa **TB infekcijom:**

- nemaju simptome
- ne osje}aju se bolesni
- ne mogu prenijeti TB bakcile na druge

TB bakcili mogu `ivjeti u tijelu godinama a da ne prouzrokuju bolest. Ako osoba koja ima infekciju ne uzima lijekove mo`e se razboliti od TB. Onda se TB bakcili mogu prenijeti na druge osobe.

Mogu}e je da imate TB infekciju ako ste ikada bili u blizini nekoga tko boluje od TB (iako niste znali da oni boluju od TB). Ljudi ne znaju da imaju TB infekciju ukoliko se ne testiraju.

Mantoux test ko`e

Najbolji na~in da se ustanovi da li imate TB infekciju je Mantoux test ko`e. Ovaj se test obi~no izvodi na ruci. Mala injekcija se upotrebljava da se ubrizga teku}ina za testiranje (zove se tuberculin) ispod ko`e. Mo`e do}i do crvenila i otoka. Trebate se javiti medicinskom osoblju 2-3 dana nakon testiranja da se ustanovi da li imate TB infekciju. Mo`da bude potrebno da se testirate ponovo da bi se ustanovilo za sigurno.

Mantoux test }e pokazati da li imate TB infekciju

Iako ste imali BCG (Bakcile Calmette-Guerin) cjepivo u pro{losti, mo`ete imati Mantoux test. BCG cjepivo ne mora uvijek prouzrokovati reakciju na Mantoux test i nije pouzdana za{tita od tuberkuloze.

Ako Mantoux test poka`e da imate TB infekciju ili ako imate simptome tuberkuloze, obavit }e se rendgenske snimke grudnog ko{a, da se vidi dali su va{a plu}a o{te}ena od tuberkuloznih klica. Da bi se utvrdilo da li }e te uzimati lijekove dodatna testiranja bi mogla biti potrebna.

Lije~enje TB infekcije

Mogu}e je da vam se preporu~i **uzimanje lijekova da bi se sprije~io razvoj tuberkuloze u budu}nosti.** Lijekovi se obi~no preporu~avaju ako:

- ste bili u blizini osobe koja ima tuberkulozu
- rendgenski snimci grudnog ko{a pokazuju da ste imali tuberkulozu u pro{losti
- imate zdravstveno stanje koje pove}ava rizik infekcije tuberkulozom

Lije~enje tuberkuloze

Lijekovi su na raspolaganju za lije~enje TB. **TB se mo`e izlije~iti ako se lijekovi uzimaju kako ih je doktor propisao.**